

Heavy metal concentrations and their possible sources in paddy soils of a modern agricultural zone, southeastern China

Chunfa Wu · Limin Zhang

Received: 12 December 2008 / Accepted: 11 April 2009 / Published online: 25 April 2009
© Springer-Verlag 2009

Abstract In a typical modern agricultural Zone of southeastern China, Haining City, 224 topsoil samples were collected from paddy fields to measure the total concentrations of copper (Cu), lead (Pb), zinc (Zn), cadmium (Cd), chromium (Cr), mercury (Hg), arsenic (As) and cobalt (Co). The total concentrations ranged from 15.30 to 78.40 mg kg⁻¹ for Cu, 20.10 to 41.40 mg kg⁻¹ for Pb, 54.98 to 224.4 mg kg⁻¹ for Zn, 0.04 to 0.24 mg kg⁻¹ for Cd, 54.90 to 197.1 mg kg⁻¹ for Cr, 0.03 to 0.61 mg kg⁻¹ for Hg, 3.44 to 15.28 mg kg⁻¹ for As, and 7.17 to 19.00 mg kg⁻¹ for Co. Chemometric techniques and geostatistics were utilized to quantify their spatial characteristics and define their possible sources. All eight metals had a moderate spatial dependency except that Pb had a strong spatial dependency. Both factor analysis and cluster analysis successfully classified the eight metals into three groups or subgroups, the first group included Cu, Zn and Cr, the second group included Cd, As and Co, and the last group included Pb and Hg. The Cu, Zn and Cr concentrations in majority samples were higher than their local background concentrations and they were highly correlated ($r > 0.80$), indicating that they had similar pollution source

and anthropic factor controlled their spatial distribution; the Cd, As and Co concentrations in majority samples were lower than their local background concentrations, indicating that the source of these elements was mainly controlled by natural factors; the mean concentration of Pb exhibited generally low level, close to its local background concentration, the Hg concentration in about half of samples was higher than its local background concentration, and they were poor correlated with the other metals, indicating that the source of Pb and Hg was common controlled by natural factor and anthropic factor.

Keywords Paddy field · Heavy metals · Spatial distribution · Source analysis

Introduction

Soil contamination, especially agricultural soil contamination had become a serious environmental problem because it poses a serious threat to human health by entering into food chains and to environmental security by leaching into ground water (Romic and Romic 2003; Shen and Chen 2000). To solve the food safety problem, modern agriculture and facilities agriculture, encouraged in densely populated developing countries such as China, and many modern agricultural demonstration zones (MADZs) were built in China from the 1980s, twentieth century. The rapid industrialization of agriculture has resulted in the incorporation of several types of pollutants to soil, such as heavy metal (HM). In numerous kinds of soil pollutions, HM contaminations have become an important environmental issue because of their nonbiodegradable nature and long biological half-live for elimination from the body (Raghunath et al. 1999; Gallego et al. 2002; Cui et al.

C. Wu
Yantai Institute of Coastal Zone Research for Sustainable
Development, Chinese Academy of Sciences,
264003 Yantai, China

C. Wu (✉)
Key Laboratory of Soil Environment and Pollution Remediation,
Institute of Soil Science, Chinese Academy of Sciences,
71 East Beijing Rd, 210008 Nanjing, China
e-mail: wchf1680@sina.com

L. Zhang
Agricultural Extension Service, 314400 Haining, China

2005). Agricultural land is fundamental resource of agricultural production that provides livelihood for the majority of people in developing countries. It is essential to identify the source of pollution and the pathway of soil contaminations enter into agricultural soil, i.e., pollution source analysis, in order to be able to formulate a policy to control or eliminate pollution and ensure food safety.

Source analysis was one of the major concerns of the present environmental study. Multivariate analysis, such as factor analysis (FA) and cluster analysis (CA) has been widely used to assist the interpretation of environmental data (e.g., Tuncer et al. 1993; Einax and Soldt 1999) and to distinguish between natural and anthropogenic inputs (e.g., Jobson 1991; Hopke 1992; Facchinelli et al. 2001; Gallego et al. 2002; Lucho-Constantino et al. 2005; Zhang 2006; Luo et al. 2007). Under the common influences of natural and anthropic input such as the utilization of chemical fertilizers and pesticides, the source of HMs in agricultural soil become more complex. Meanwhile, due to the heterogeneity of soil itself, concentrations of HMs vary remarkably over space (Luo et al. 2007). Only relying on multivariate statistics or geostatistics, it is very difficult both to disclose their sources and to characterize their regional variations. Fortunately, a combination of these two methods provides an appropriate solution and has been proved to be feasible in some studies (Chen et al. 2008; Facchinelli et al. 2001).

In the past decades, a large amount of soil pollution surveys on HMs have been carried out at different scales and there were many studies reported in the scientific literatures (Benvenuti et al. 1995; Chen et al. 2008; Cullbard et al. 1988; Li et al. 2004; Lucho-Constantino et al. 2005; Micó et al. 2008; Shi et al. 2007; von Steiger et al. 1996). However, few investigations have been conducted in modern agricultural zone of developing countries, where the agriculture soil ecosystem usually has higher pollution risk for agricultural industrialization. In this study, we choose one of typical MADZs in Southeastern China, Haining City, which was surrounded by rapidly industrialized cities such as Shanghai City, Hangzhou City and Suzhou City. Previous studies showed that land use/cover affected the HMs accumulation and spatial distribution, and the HM sources in different land use/cover are always different (Bloemen et al. 1995; Kelly et al. 1996; Rui et al. 2008). To remove the effect of land use/cover on HMs accumulation, we only study HMs in the main land use/cover -paddy fields. Paddy field is widely distributed in the southeast region of China. The general objectives of this study were to (1) investigate the total concentrations of HMs in topsoil of paddy fields, (2) characterize their spatial distribution, (3) define their possible sources based on data analysis with the use of chemometric techniques and geostatistics.

Materials

Descriptions of study area

The study area is located in the Hang-Jia-Hu Plain, northeastern region of Zhejiang Province, China (Fig. 1). The study area is bounded by east longitude 120°18'–120°52' and north latitude 30°15'–30°35' with a total area of about 731 km². The study area is in the northern subtropical zone of monsoonal climate with a temperate and humid climate throughout the year with four distinct seasons. The average annual temperature is 15.9°C and the mean annual precipitation is approximately 1,190 mm. Paddy field is the dominant land use/land cover of arable land and paddy soil (Gleysols), one kind of anthropogenic soil that is the main soil type, although there are several other soil types in the area.

Sampling design and soil analysis

A total of 224 soil samples (0–15 cm) were collected from paddy fields in November 2005 with consideration of land use uniformity and soil types to ensure all samples were located in paddy fields and a soil sample was collected from each soil type (Fig. 1). When sampling, soils in top layer of 6–8 points in each site of an area of about 0.1–0.2 ha were collected and then fully mixed, and finally divided into parts of 1–2 kg each. Only one of the parts was packed with a bag and brought back laboratory for analysis. All sample sites were recorded using a hand-held global position system (GPS). All samples were air-dried at room




Fig. 1 The location of study area and distribution of sampling points

temperature (20–22°C), removed stones or other debris, and then sieved to 2 mm polyethylene sieve. Portions of soil samples (about 100 g) were ground in an agate grinder and sieved through 0.149 mm mesh. The prepared soil samples were then stored in polyethylene bottles for analysis.

Total metal concentration is useful in identifying the pollution source and the potential for contamination (He et al. 1997; Jung 2001). In this study, we analyzed soil pH, soil organic matter concentration (SOM) and the total concentrations of copper (Cu), lead (Pb), zinc (Zn), cadmium (Cd), chromium (Cr), mercury (Hg), cobalt (Co) and arsenic (As). In this study, As was also seemed as one of HMs for it has similar properties as HMs. Soil pH was measured by pH meter with a soil/water ratio of 1:2.5. SOM was determined by wet oxidation at 180°C with a mixture of potassium dichromate and sulfuric acid (Agricultural Chemistry Committee of China 1983). Before the concentrations of HM were measured, soil samples had been digested with a mixture of nitric acid (HNO₃) and perchloric acid (HClO₄) (Agricultural Chemistry Committee of China 1983). Total concentrations of Cu, Pb, Cd and Co were measured by inductively coupled plasma mass spectrometry (ICP-MS). Calibration of ICP-MS was carried out using standard solutions in the concentrations ranging from 10 to 1,000 µg l⁻¹. For analyte concentrations outside the calibration range, dilution in 1% HNO₃ was used. Zn and Cr were analyzed by inductively coupled plasma optical emission spectrometry (ICP-OES). In addition, the concentrations of As and Hg in the digested solution were determined by hydrid generation atomic fluorescence spectrometry (HG-AFS) and cold vapor atomic fluorescence spectrometry (CV-AFS), respectively.

The validity of the whole analyzing procedure was checked using the certified reference materials (CRM) IEAE-Soil-7 and CRM 277. Analyses of CRMs, replicate samples and blanks were performed after every ten samples and were carried through the entire sample preparation and analytical process. The precision of the measurements, estimated by carrying out 22 replicates, was in the range of 2.1–4.3% RSD (relative standard deviation). The average recovery rates of certified elements were between 93 and 105%. Method detection limit (MDL) for each heavy metal was Cu, As and Co, 1 µg g⁻¹; Pb and Zn, 2 µg g⁻¹; Cr, 5 µg g⁻¹; Cd, 20 ng g⁻¹; Hg, 2 ng g⁻¹, respectively.

Methods

Data transformation

In multivariate statistics and linear geostatistics, a normal distribution for the variables under study was desirable

(Webster and Oliver 2001; Gallego et al. 2002; Zhang and McGrath 2004). To avoid result distortions and low levels of significance, data transformation has been performed on all measured values. In numerous data transformation methods, logarithmic transformation is widely applied (Webster and Oliver 2001). However, it is often found that environmental variables do not always follow the lognormal distribution (Zhang and McGrath 2004). In our study, the Box–Cox transformation was used to make the data more normal and less skewness. The detail theory and process of Box–Cox transformation can be found in textbooks and monographs such as Box and Cox (1964).

Geostatistical analysis

Geostatistical estimation allows one to predict values at unsampled locations by taking into account of the spatial correlation between estimated and sampled points (i.e., spatial variability) and minimizes the variance of estimation error (Saito et al. 2005). Many studies used geostatistics to describe the spatial distribution of pollutants in contaminated soils (Atteia et al. 1994; Arrouays et al. 1996; Carlon et al. 2001; Goovaerts 1997; Meuli et al. 1998). The semivariogram, main components of kriging, is effective tools for evaluating spatial variability (Boyer et al. 1991; Cahn et al. 1994), which provide a clear description of the spatial structure of variable and some insight into possible processes affecting its distribution (Paz González et al. 2001; Wang and Tao 1998; Webster and Oliver 1990). In this study, anisotropy of variograms was not found for the data set; all the semivariograms in isotropic form were fit using a spherical model, exponential model, Gaussian model and linear model, respectively, and the best-fit model of them was applied to kriging interpolation. Ordinary kriging was chosen to create the spatial distribution maps of HM after data transformation, using the nearest 16 sampling points and a maximum searching distance equal to the range distance of the variable. For a more technical description of kriging and the semivariogram (see Cressie 1993 or Webster and Oliver 2001). Software version 7.0 of GS+ Geostatistics for the Environmental Sciences was used to perform all the geostatistical computations.

Multivariate analysis

Multivariate statistical solutions are mathematical hypotheses and their interpretation requires environmental knowledge. These techniques have been widely used to assist the interpretation of environmental data and to distinguish between natural and anthropogenic inputs of HMs (e.g., Dudka 1992; Einax and Soldt 1999; Luch-

Constantino et al. 2005). In this study, multivariate analysis (correlation analysis, FA and CA) was interpreted to build different groups of HMs with approximately the same or similar source.

Inter element relationships can provide interesting information on HM sources and pathways (Manta et al. 2002). In the study area, correlation analysis was applied to identify the relationships between total concentration of HMs and soil properties (soil pH and SOM). Also, correlation analysis was performed considering total concentration of HMs to check relationships among them. Spearman nonparametric correlation coefficient was used for most of variables in this study positively skewed. FA is an useful statistics tool that can extract latent information from multidimensional data and group the measured elements into fewer groups (Gallego et al. 2002). The original data matrix is decomposed into the products of a matrix of factor loadings and a matrix of factor scores plus a residual (Kowalkowski et al. 2006). In our study, the common factors were extracted by principal components method to gain the observed correlation matrix. Furthermore, in order to facilitate the interpretation of results, varimaxrotation was applied because orthogonal rotation minimized the number of variables with a high loading (Gallego et al. 2002). CA is often coupled to FA to confirm results and provide grouping of variables (Facchinelli et al. 2001). In our study, CA was performed according to the between-groups linkage method. Results are shown in a dendrogram, where steps in the hierarchical clustering solution and values of the distances between clusters are based on correlation coefficients (Pearson coefficient). Dendrogram was undertaken according to the average linkage clustering that calculates dissimilarity between clusters considering the cluster average values. Multivariate analysis was performed using SPSS R13.0 for Windows.

Results and discussion

Descriptive statistics

The soil pH in the area was in range of 4.44–8.30 with a mean of 6.57 and the SOM was in range of 5.7–45.9 g kg⁻¹ with a mean of 22.43 g kg⁻¹. A descriptive summary of soil HMs (Cu, Pb, Zn, Cd, Cr, Hg, As and Co) was listed in Table 1. The results showed that the concentrations of all elements had wide range, suggesting extrinsic factors affect these element concentrations in the topsoil of paddy fields. The mean concentrations of Cu, Pb, Zn, Cr and Hg were 27.87, 29.59, 87.30, 77.60 and 0.17 mg kg⁻¹, respectively, higher than their background values at Zhejiang Province scale (Zhejiang Soil Survey Office 1994; Cheng et al. 2006). The mean Cr (87.30 mg kg⁻¹) and Hg (0.17 mg kg⁻¹) concentrations of paddy fields in study area were also significantly higher than the mean Cr (67.29 mg kg⁻¹) and Hg (0.13 mg kg⁻¹) of agricultural soils in Zhejiang Province (Cheng et al. 2007), respectively, suggesting the Cr and Hg enrichments in study area were more serious than that in the other area of Zhejiang Province. In the study area, the Cu, Zn and Cr concentrations in more than 85% samples were higher than their local background concentrations and their mean concentrations were significantly higher than their local background concentration, showed Cu, Zn and Cr were moderate high enriching in majority area; the Cd, As and Co concentrations in more than 65% samples were lower than their local background concentrations, showed Cd, As and Co were enriching in minority area; the Pb concentration in more than 90% samples was higher than its local background concentration and its mean concentration was slight higher than its local background concentration, showed that Pb was slight enriching in majority area; the Hg concentration in about half samples exceeding it

Table 1 Summary statistics for the eight heavy metal concentrations in paddy soils (mg kg⁻¹) (*n* = 224)

	Cu	Pb	Zn	Cd	Cr	Hg	As	Co
Minimum	15.30	20.10	54.98	0.04	54.90	0.03	3.44	7.17
Maximum	78.40	41.40	224.4	0.24	197.1	0.61	15.28	19.00
Mean	27.87	29.59	87.30	0.14	77.60	0.17	8.32	13.02
Median	27.80	29.40	86.15	0.13	77.10	0.15	8.01	12.94
CV(%)	26.0	11.7	20.5	27.2	17.8	49.8	28.8	17.4
Skewness	2.20	0.34	2.13	0.33	3.52	1.62	0.44	0.10
Kurtosis	11.14	0.84	13.95	0.07	25.77	3.91	-0.06	-0.59
Background value ^a	17.76	25.61	69.00	0.17	54.60	0.15	9.33	14.00
Percentage ^b	96.9%	90.2%	85.3%	17.0%	100%	49.1%	34.8%	32.6%

^a Soil heavy metal background value of Zhejiang Province (Zhejiang Soil Survey Office 1994; Cheng et al. 2006)

^b The percent of soil heavy metal exceeding the background value of Zhejiang Province

background value, showed that the topsoil was partial enriching by Hg. The enrichment of Pb probably due to the exhaust emissions from vehicle for the amount of vehicle sharply increase in recent years. The coefficient of variation (CV) of the metals was calculated as indicator of heterogeneity, which varied from 11.7% for Pb to 49.8% for Hg. For most metal concentrations, the CVs exceeded 20% indicating considerable variability. High concentrations of Hg coupled with high coefficients of variation suggest anthropogenic sources for Hg (Manta et al. 2002).

Compared to enriched elements in topsoil, the mean concentrations of Cd, As and Co exhibited generally low levels, close to their local background concentrations. The mean Cd concentration (0.14 mg kg⁻¹) of paddy fields in study area was significantly lower than that (0.20 mg kg⁻¹) of agricultural soils in Zhejiang Province (Cheng et al. 2007). There were 38 (17.0%) soil samples for Cd, 78(34.8%) for As, and 73(32.6%) for Co exceeding the background values at Zhejiang Province scale, respectively. It indicated that Cd, As and Co enrichment did not exist in the topsoil of majority area, and natural factor as an important role for controlling their distribution.

Data transformation

Prior to multivariate analysis and spatial analysis, the normality of all element concentrations was checked. The parameters of skewness, kurtosis, and the significance level of Shapiro–Wilk’s test for normality (S–K *p*) were shown in Table 2. In our study, it was found that only Cd and Co passed the Shapiro–Wilk’s normality test (S–K *P* > 0.05) before data transformation, whereas other variables were all strongly skewed, with skewnesses greater than 0. Their kurtoses were also very sharp, caused by the fact that the majority of samples were clustered at relatively low values.

Compared with log-transformation, the Box–Cox transformation has significantly reduced the skewness values of data sets, pushing them toward “0” although there were four transformed variables did not pass the Shapiro–Wilk’s normality test.

Geostatistical analysis

Spatial structure analysis

Soil HMs in environment science have spatial structures, including spatial autocorrelation. In this study, the geostatistics method was used to spatial structure analysis. The semivariogram and the fitted models for 8 metals were presented in Fig. 2. The parameters of semivariogram for each element were summarized in Table 3. Semivariograms showed that soil Cu, Pb, Zn, Cd and Cr were all fitted for an exponential model, whereas soil Hg, As and Co were all best fitted for a spherical model.

The Nugget/Sill ratio was assumed to be a criterion to classify the spatial dependence of HMs. The ratio values lower than 25% and higher than 75% corresponded to strong and weak spatial dependency, respectively, while the ratio values between 25 and 75% corresponded to moderate spatial dependence (Cambardella et al. 1994). Usually, strong spatial dependence of soil properties can be attributed to intrinsic factors, and weak spatial dependence can be attributed to extrinsic factors (Cambardella et al. 1994). The Nugget/Sill ratios of all models lower than 25% and higher than 75% except for the model for Pb lower than 25%. It indicated that all metals had a moderate spatial dependency except that Pb had a strong spatial dependency, and the spatial dependence of Pb in study area may be attributed to natural factor.

Range value is a measure of extension where autocorrelation exists (Li et al. 2007; Webster and Oliver

Table 2 Skewness, kurtosis, and significance level of Shapiro–Wilk’s test for normality (S–W *p*) of the raw, log-transformed (Log), and Box–Cox transformed (Box–Cox) data sets of eight soil heavy metals (*n* = 224)

Data set	Parameter	Cu	Pb	Zn	Cd	Cr	Hg	As	Co
Raw data	Skewness	2.20	0.34	2.13	0.33	3.52	1.62	0.44	0.10
	Kurtosis	11.14	0.84	13.95	0.07	25.77	3.91	-0.06	-0.59
	S–W <i>p</i>	0.01	0.03	0.01	0.07	0.01	0.01	0.02	0.10
Log	Skewness	0.51	-0.12	0.41	-0.63	1.30	-0.17	-0.29	-0.30
	Kurtosis	1.62	0.68	1.96	1.12	6.41	1.08	-0.24	-0.27
	S–W <i>p</i>	0.01	0.10	0.01	0.01	0.01	0.01	0.10	0.05
Box–Cox	Skewness	0.39	-0.12	0.04	-0.11	0.37	-0.17	-0.29	-0.30
	Kurtosis	0.13	0.68	0.54	0.19	-0.07	1.08	-0.24	-0.27
	S–W <i>p</i>	0.01	0.10	0.01	0.10	0.01	0.01	0.10	0.05
λ		-1.00	0.00	-0.50	0.50	-2.00	0.00	0.00	0.00

S–W *p* is the significance level of Shapiro–Wilk’s test for normality; λ is the parameter of the Box–Cox transformation method




Fig. 2 Experimental semivariograms of soil heavy metals with fitted models

1990). The ranges of spatial autocorrelation varied greatly from metal to metal. The variability of Pb was dominated by relatively short-range spatial correlation (3, 270 m),

whereas the spatial correlation structures of the other elements had long ranges (16,500–60,090 m). Short-range spatial correlation of Pb, suggesting the anthropogenic

Table 3 Best-fitted semivariogram models of soil heavy metals and their parameters

Soil elements	Model	C_0	$C_0 / (C + C_0)$	Range (m)	R^2
Cu	Exponential	2.6E-5	0.303	39150	0.879
Pb	Exponential	9.5E-4	0.073	3270	0.771
Zn	Exponential	3.9E-5	0.382	16500	0.846
Cd	Exponential	9.2E-4	0.497	60090	0.900
Cr	Exponential	8.1E-10	0.291	31230	0.915
Hg	Spherical	1.5E-1	0.498	39050	0.907
As	Spherical	4.2E-2	0.425	27200	0.977
Co	Spherical	1.3 E-2	0.331	34190	0.978

C_0 , nugget variance, and $C + C_0$, sill variance

factor affecting Pb distribution, however, low Nugget/Sill ratio (0.073) of Pb, with low CV (11.7%), suggesting the natural factor also affecting Pb distribution. Based on above analysis, it is reasonable to conclude that Pb in topsoil of paddy fields was controlled by both natural factor and anthropic factor. Comparing the range of eight elements, it was found that Cd has longer effective range than Cu, Pb, Zn, Cr, As and Hg, indicating that the Cd has better spatial structure and less variation caused by extrinsic factors. According to the result of descriptive statistics, the Cd concentration of majority samples was lower than the local background concentration, suggesting the natural factor as an important role controlling Cd distributions.

Spatial distribution

In order to know distribution patterns of eight elements, kriging interpolation was used to obtain the filled contours maps (Fig. 3). The spatial maps of six elements (Cu, Zn, Cd, Cr, As and Co) showed similar distribution trends with the high concentrations in the northeast area and low concentrations in the southwest area. According to spatial maps, the Hg concentration in majority area was lower than the local background value, and only southwest area had high Hg, suggesting the main pollution source in southwest area was different from that in the other unpolluted area. Southwestern of study area adjoins the traditional industry area of Xiaoshan City, industrial fumes and atmosphere deposition could also be an important source of Hg enrichment in this area as the result some previous studies (Wang et al. 2003; Mukherjee and Zevenhoven 2006). The spatial maps of Pb showed local spatial variability and its distribution pattern seemed more irregular than the other elements, suggesting the anthropogenic factor as an important role affecting Pb distributions.

Multivariate statistical analysis

Correlation analysis

To obtain the valuable information, the Spearman non-parametric correlation coefficients were calculated between eight elements and related soil properties (pH and Organic matter), and the results were shown in Table 4. SOM showed good correlation with Cu, Pb, Zn, Cr and As, indicating the strong adsorption effect of organic matter on these elements. All elements concentrations had significant correlation with pH except Cd, Hg and Co, however, the correlation coefficients were relatively low. It indicated that the effect of pH on the trace metals distribution was relatively limited. Strong correlations were found between Cu, Zn and Cr, indicating the main sources of these elements were similar or same. However, Hg was relatively poor correlated with the Cu, Zn and Cd, indicating the main source of Hg was different from the Cu, Zn and Cr.

Factor analysis

To reduce the high dimensionality of variable space and better understand the relationships among HMs, FA was applied to the transformed data matrixes. The results of FA for HM concentrations were presented in Tables 5 and 6. According to the results of the initial eigenvalues in Table 5, two factors were extracted from the available data set, which account for over 76% of the total variation of soil elements (Cu, Pb, Zn, Cd, Cr, Hg, As and Co). In Table 6, the rotated component matrix for data indicated that Cu, Zn, Cr, As and Co were strongly associated with the first component (F1) with similar high absolute values. Soil Cd also associated with the first component with moderate absolute value. Elements Pb and Hg were shown in the second component (F2) with high values.

F1 was responsible for 57.8% of total variance and was dominated by Cu, Zn, Cr, As, Co and Cd. Cu, Zn and Cr were strong positively associated in F1, and As, Co and Cd were negatively associated in F1, suggesting Cu, Zn and Cr may have similar source and As, Co and Cd also may have similar source, however, they was different. In our study, it was found that the mean concentrations of Cu, Zn and Cr exhibited high lever, the concentrations of exceeding 85% samples higher than their local background concentrations, suggesting anthropic factor controlling their distribution; the mean concentrations of As, Co and Cd exhibited generally low levels, lower than their local background concentrations, suggesting natural factor controlling their distribution. F2 explained 18.2% of total variance and was mainly attributed to Pb and Hg. The mean concentrations of Pb and Hg in study area were slight higher than local




Fig. 3 The prediction maps of soil heavy metals

background values, suggesting both anthropic factor and natural factor controlling their distribution.

Cluster analysis

In order to discriminate distinct groups of HMs as tracers of natural or anthropic source, CA was performed on eight elements. The results were illustrated with the dendrogram (Fig. 4) which enabled the identification of two main groups to describe the complex reality of the area studied,

distinguishing Cu, Zn and Cr (Group I) from As, Co, Cd, Hg and Pb (Group II).

Group I was made up of those elements which involved in F1 that were strong positively associated, seem to be controlled by anthropic factor. Several studies have shown that soil acts as a long-term sink for Cu and Zn when it is fertilized with manure and chemical fertilizers (L'Herroux et al. 1997; Chang and Page 2000; Nicholson et al. 2003). Overuse or misuse of fertilizers and manure was popular existed in the area, and these elements were enriching in

Table 4 The nonparametric correlations between the contents of eight heavy metals and two selected soil properties ($n = 224$)

	Cu	Pb	Zn	Cd	Cr	Hg	As	Co
Cu	1							
Pb	0.63**	1						
Zn	0.88**	0.56**	1					
Cd	0.48**	0.28**	0.52**	1				
Cr	0.84**	0.48**	0.81**	0.48**	1			
Hg	-0.06	0.41**	-0.10	-0.02	-0.21**	1		
As	0.75**	0.42**	0.73**	0.45**	0.71**	-0.18**	1	
Co	0.72**	0.35**	0.71**	0.50**	0.80**	-0.31**	0.65**	1
Soil pH	-0.30**	-0.19**	-0.20**	0.05	-0.22**	-0.08	-0.25**	-0.06
SOM	0.64**	0.48**	0.59**	0.37**	0.49**	0.06	0.59**	0.33**

*Significant at the 0.05 probability level (two-tailed)

**Significant at the 0.01 probability level (two-tailed)

Table 5 Total variance explained for heavy metal concentrations by principal component analysis

Component	Initial eigenvalues			Extraction sums of squared loadings			Rotation sums of squared loadings		
	Total	% Of variance	Cumulative (%)	Total	% Of variance	Cumulative (%)	Total	% Of variance	Cumulative (%)
1	4.625	57.81	57.81	4.625	57.81	57.81	4.591	57.39	57.39
2	1.454	18.17	75.98	1.454	18.17	75.98	1.488	18.60	75.98
3	0.677	8.46	84.44						
4	0.421	5.27	89.71						
5	0.314	3.93	93.64						
6	0.234	2.93	96.57						
7	0.176	2.20	98.77						
8	0.098	1.23	100.00						

Table 6 Component matrix and rotated component matrix for heavy metal concentrations

Element	Component matrix		Rotated component matrix	
	F1	F2	F1	F2
Cu	0.937	-0.095	0.922	-0.191
Pb	-0.585	0.686	-0.511	0.742
Zn	0.914	-0.036	0.906	-0.131
Cd	-0.641	-0.013	-0.639	0.053
Cr	0.903	0.102	0.908	0.008
Hg	0.136	0.936	0.232	0.917
As	-0.799	-0.076	-0.803	0.007
Co	-0.828	-0.285	-0.853	-0.198

Extraction method: principal component analysis

Rotation method: varimax with Kaiser normalization

the topsoil of majority area. It indicated that fertilization may be the main pollution source of Cu, Zn and Cr.

In Group II, Hg and Pb were the extraordinary contamination elements and their origin were not entirely the same as Cd, As and Co, so their cluster distance were the

furthest from other three elements (Cd, As and Co). It is reasonable to divided the Group II into two subgroups in pollution source identification; the first subgroup (Group II₁) including As, Co and Cd and the second subgroup (Group II₂) including Hg and Pb. Group II₁ was made up of those elements which involved in F1 that were negatively associated. This was in agreement with factor analysis results Group II₂ was made up of those elements which involved in F2, and they were poor correlated with the metals of Group II₁ and Group I, suggesting the main source of Pb and Hg was different from that of Group II₁ (natural factors) and Group I (anthropic factor), both natural factors and anthropic factor common controlling Pb and Hg distribution. And this was also in agreement with factor analysis results. However, multivariate statistical methods can only be used to discriminate natural or anthropic sources of HMs but cannot quantify and ascertain their concrete sources.

Based on above analysis, it is reasonable to conclude that Cu, Zn and Cr in topsoil of paddy fields are mainly controlled by anthropic inputs, Hg and Pb in topsoil of paddy fields are controlled by both anthropic inputs and




Fig. 4 Dendrogram with between-groups linkage method of the cluster analysis of soil heavy metals

parent materials, whereas As, Co, and Cd in topsoil of paddy fields appear to be mainly associated with parent materials.

Conclusions

In the study area, Cu, Zn and Cr were moderate high enriching in majority area, Pb was slight enriching in majority area, Hg was enriching in partial area and Cd, As and Co were enriching in minority area. Cu, Zn, Cd, Cr, As, Hg and Co had a moderate spatial dependency, and the soil Pb had a strong spatial dependency. The Cu, Zn, Cd, Cr, As and Co in topsoil of paddy fields showed similar distribution trends with the high concentrations in the northeast area and low concentrations in the southwest area; the Hg showed a distribution trend with the high concentration in southwest area and lower concentration in the other area; and the Pb showed a distribution trend with the lower concentration in the middle area and high concentration in the other area.

The degrees of the eight HMs enrichment in topsoil of paddy fields were different in study area for the influence of anthropogenic activity on them was different. The result of chemometric techniques and geostatistics analyses indicated that the Cu, Zn and Cr were mainly controlled by anthropic factor, Cd, As and Co mainly controlled by natural factors and Pb and Hg controlled by both natural and anthropic factors.

The survey data show that over 85% of the study area existed Cu, Zn and Cr enrichment, indicating the

requirement of modifications of agricultural cultivation practices in these areas. The spatial map showed that the southwest area had high Hg, indicating the requirements of controlling Hg enrichment in this area.

Acknowledgments The authors are grateful to the grants by the National Basic Research Priorities Program (973 Program) (2002CB410810). We appreciate all the colleagues who collected and analyzed the soil samples. We also extend great appreciation to Dr. Jiaping Wu and Dr. Huiqing Huang of the Zhejiang University, for their aids in auxiliary data collection.

References

- Agricultural Chemistry Committee of China (1983) Conventional methods of soil and agricultural chemistry analysis (in Chinese). Science Press, Beijing, pp 70–165
- Arrouays D, Mench M, Amans V, Gomez A (1996) Short-range variability of fallout Pb in a contaminated soil. *Can J Soil Sci* 76:73–81
- Atteia O, Dubois JP, Webster R (1994) Geostatistical analysis of soil contamination in the Swiss Jura. *Environ Pollut* 86:315–327. doi:10.1016/0269-7491(94)90172-4
- Benvenuti M, Mascaro I, Corsini F, Lattanzi P, Parrini P, Tanelli G (1995) Mine waste dumps and heavy metal pollution in abandoned mining district of Boccheggiano southern Tuscany, Italy. *Environ Geol* 30:238–243. doi:10.1007/s002540050152
- Bloemen ML, Markert B, Lieth H (1995) The distribution of Cd, Cu, Pb, and Zn in topsoils of Osaruck in relation to land uses. *Sci Total Environ* 166:137–148. doi:10.1016/0048-9697(95)04520-B
- Box GEP, Cox DR (1964) An analysis of transformations. *J R Stat Soc B* 26:1–78
- Boyer DG, Wright RJ, Feldhake CM, Bligh DP (1991) Soil spatial variability in steeply sloping acid soil environment. *Soil Sci* 161:278–287. doi:10.1097/00010694-199605000-00003
- Cahn MD, Hummel JW, Brouer BH (1994) Spatial analysis of soil fertility for site-specific crop management. *Soil Sci Soc Am J* 58:1240–1248
- Cambardella CA, Moorman TB, Nacak JM, Parkin TB, Karlen DL, Turco RF, Konopka AE (1994) Field-scale variability of soil properties in central Iowa soils. *Soil Sci Soc Am J* 58:1501–1511
- Carlson C, Critto A, Marcomini A, Nathanail P (2001) Risk based characterisation of contaminated industrial site using multivariate and geostatistical tools. *Environ Pollut* 111:417–427. doi:10.1016/S0269-7491(00)00089-0
- Chang AC, Page AL (2000) Trace elements slowly accumulating, depleting in soils. *Calif Agr* 54(2):49–55
- Chen T, Liu XM, Zhu MZ, Zhao KL, Wu JJ, Xu JM, Huang PM (2008) Identification of trace element sources and associated risk assessment in vegetable soils of the urban-rural transitional area of Hangzhou, China. *Environ Pollut* 151:67–78. doi:10.1016/j.envpol.2007.03.004
- Cheng JL, Shi Z, Zhu YW, Liu C, Li HY (2006) Differential characteristics and appraisal of heavy metals in agricultural soils of Zhejiang Province (in Chinese). *J Soil Water Conserv* 20(1): 103–107
- Cheng JL, Shi Z, Zhu YW (2007) Assessment and mapping of environmental quality in agricultural soils of Zhejiang Province, China. *J Environ Sci China* 19:50–54. doi:10.1016/S1001-0742(07)60008-4
- Cressie N (1993) *Statistics for spatial data* (revised edition). Wiley, New York

- Cui Y, Zhu YG, Zhai R, Huang Y, Qiu Y, Liang J (2005) Exposure to metal mixtures and human health impacts in a contaminated area in Nanning, China. *Environ Int* 31:784–790. doi:10.1016/j.envint.2005.05.025
- Cullbard EB, Thornton I, Wheatley M, Moorcroft S, Thompson M (1988) Metal contamination in British urban dusts and soils. *J Environ Qual* 17(2):226–234
- Dudka S (1992) Factor analysis of total element concentrations in surface soils of Poland. *Sci Total Environ* 121:39–52. doi:10.1016/0048-9697(92)90305-C
- Einax JW, Soldt U (1999) Geostatistical and multivariate statistical methods for the assessment of polluted soils: merits and limitations. *Chemometr Intell Lab* 46:79–91. doi:10.1016/S0169-7439(98)00152-X
- Facchinelli A, Sacchi E, Mallen L (2001) Multivariate statistical and GIS based approach to identify heavy metal sources in soils. *Environ Pollut* 114:313–324. doi:10.1016/S0269-7491(00)00243-8
- Gallego JLR, Ordóñez A, Loredó J (2002) Investigation of trace element sources from an industrialized area (Avilés, northern Spain) using multivariate statistical methods. *Environ Int* 27:589–596. doi:10.1016/S0160-4120(01)00115-5
- Goovaerts P (1997) *Geostatistics for natural resources evaluation*. Oxford University Press, New York
- He M, Wang Z, Tang H (1997) Spatial and temporal patterns of acidity and heavy metals in predicting the potential for ecological impact on the Le An river polluted by acid mine drainage. *Sci Total Environ* 206:67–77. doi:10.1016/S0048-9697(97)00217-9
- Hopke PK (1992) Factor and correlation analysis of multivariate environmental data. In: Hewitt CN (ed) *Methods of environmental data analysis*. Elsevier Applied Science, London, pp 139–180
- Jobson JD (1991) *Applied multivariate data analysis*. Springer, New York
- Jung MC (2001) Heavy metal contamination of soils and waters in and around the Imcheon Au–Ag mine, Korea. *Appl Geochem* 16:1369–1375. doi:10.1016/S0883-2927(01)00040-3
- Kelly J, Thornton I, Simpson PR (1996) Urban geochemistry: a study of the influence of anthropogenic activity on the heavy metal content of soil in traditionally industrial and no-industrial areas of Britain. *Appl Geochem* 11:363–370. doi:10.1016/0883-2927(95)00084-4
- Kowalkowski T, Zbytniewski R, Szpejna J, Buszewski B (2006) Application of chemometrics in river water classification. *Water Res* 40:744–752. doi:10.1016/j.watres.2005.11.042
- L'Herroux L, Le Roux S, Appriou P, Martinez J (1997) Behaviour of metals following intensive pig slurry applications to a natural field treatment process in Brittany France. *Environ Pollut* 97:119–130. doi:10.1016/S0269-7491(97)00072-9
- Li X, Lee SL, Wong SC, Shi W, Thornton I (2004) The study of metal contamination in urban soils of Hong Kong using a GIS-based approach. *Environ Pollut* 129:113–124. doi:10.1016/j.envpol.2003.09.030
- Li BG, Ran Y, Cao J, Liu WX, Shen WR, Wang XJ, Coveney RM, Tao S (2007) Spatial structure analysis and kriging of dichlorodiphenyltrichloroethane residues in topsoil from Tianjin, China. *Geoderma* 141(1–2):71–77. doi:10.1016/j.geoderma.2007.05.004
- Lucho-Constantino CA, Álvarez-Suárez M, Beltrán-Hernández RI, Prieto-García F, Poggi-Valardo HM (2005) A multivariate analysis of the accumulation and fractionation of major and trace elements in agricultural soils in Hidalgo State, Mexico irrigated with raw wastewater. *Environ Int* 31:313–323. doi:10.1016/j.envint.2004.08.002
- Luo W, Wang T, Lu Y, Giesy JP, Shi Y, Zheng Y, Xing Y, Wu G (2007) Landscape ecology of the Guanting Reservoir, Beijing, China: multivariate and geostatistical analyses of metals in soils. *Environ Pollut* 146:567–576. doi:10.1016/j.envpol.2006.08.001
- Manta DS, Angelone M, Bellanca A, Neri R, Sprovieri M (2002) Heavy metals in urban soils: a case study from the city of Palermo (Sicily), Italy. *Sci Total Environ* 300:229–243. doi:10.1016/S0048-9697(02)00273-5
- Meuli R, Schulin R, Webster R (1998) Experience with the replication of regional survey of soil pollution. *Environ Pollut* 101:311–320. doi:10.1016/S0269-7491(98)00064-5
- Micó C, Recatalá L, Peris M, Sánchez J (2008) Discrimination of lithogenic and anthropogenic metals in calcareous agricultural soils: a case study of the Lower Vinalopó Region (SE Spain). *Soil Sediment Contam* 17:467–485. doi:10.1080/15320380802304367
- Mukherjee AB, Zevenhoven R (2006) Mercury in coal ash and its fate in the Indian subcontinent: a synoptic review. *Sci Total Environ* 368:384–392. doi:10.1016/j.scitotenv.2005.08.022
- Nicholson FA, Smith SR, Alloway BJ, Smith CC, Chambers BJ (2003) An inventory of heavy metal input to agricultural soil in England and Wales. *Sci Total Environ* 311(1–3):205–219. doi:10.1016/S0048-9697(03)00139-6
- Paz González A, Taboada Castro MT, Vieira SR (2001) Geostatistical analysis of heavy metals in a one-hectare plot under natural vegetation in a serpentine area. *Can J Soil Sci* 81(4):469–479
- Raghunath R, Tripathi RM, Kumar AV, Sathe AP, Khandekar RN, Nambi KSV (1999) Assessment of Pb, Cd, Cu, and Zn exposures of 6 to 10-year-old children in Mumbai. *Environ Res* 80:215–221. doi:10.1006/enrs.1998.3919
- Romic M, Romic D (2003) Heavy metals distribution in agricultural topsoils in urban area. *Environ Pollut* 43:795–805
- Rui YK, Qu LC, Kong XB (2008) Effects of soil use along Yellow River Basin on the pollution of soil by heavy metals (in Chinese). *Spectrosc Spect Anal* 28(4):934–936
- Saito H, McKenna A, Zimmerman DA, Coburn TC (2005) Geostatistical interpolation of object counts collected from multiple strip transects: ordinary Kriging versus finite domain kriging. *Stoch Environ Res Risk Assess* 19:71–85. doi:10.1007/s00477-004-0207-3
- Shen Z, Chen HM (2000) Bioremediation of heavy metal polluted soils. *Rural Eco Environ* 16(2):39–44
- Shi JC, Wang HZ, Xu JM, Wu JJ, Liu XM, Zhu HP, Yu CL (2007) Spatial distribution of heavy metals in soils: a case study of Changxing, China. *Environ Geol* 52:1–10. doi:10.1007/s00254-006-0443-6
- Tuncer GT, Tuncel SG, Tuncel G, Balkas TI (1993) Metal pollution in the Golden Horn, Turkey—contribution of natural and anthropogenic components since 1913. *Water Sci Technol* 28:59–64
- von Steiger B, Webster R, Schulin R, Lehmann R (1996) Mapping heavy metals in polluted soil by disjunctive kriging. *Environ Pollut* 94:205–215. doi:10.1016/S0269-7491(96)00060-7
- Wang XJ, Tao S (1998) Spatial structures and relations of heavy metal content in wastewater irrigated agricultural soil of Beijing's eastern farming regions. *B Environ Contam Tox* 61:261–268. doi:10.1007/s001289900757
- Wang D, Shi X, Wei S (2003) Accumulation and transformation of atmospheric mercury in soil. *Sci Total Environ* 304:209–214. doi:10.1016/S0048-9697(02)00569-7
- Webster R, Oliver MA (1990) *Statistical methods in soil and land resource survey*. Oxford University Press, London
- Webster R, Oliver MA (2001) *Geostatistics for environmental scientists*. Wiley, Chichester, pp 37–103

- Zhang CS (2006) Using multivariate analyses and GIS to identify pollutants and their spatial patterns in urban soils in Galway, Ireland. *Environ Pollut* 142:501–511. doi:10.1016/j.envpol.2005.10.028
- Zhang C, McGrath D (2004) Geostatistical and GIS analyses on soil organic carbon concentrations in grassland of southeastern Ireland from two different periods. *Geoderma* 119:261–275. doi:10.1016/j.geoderma.2003.08.004
- Zhejiang Soil Survey Office (1994) Zhejiang soils (in Chinese). Zhejiang Technology Press, Hangzhou, pp 556–562